

# Beginning of the End or the End of the Beginning?

Opinion

## COVID-19 in 2022—The

**Carlos del Rio, MD** Division of Infectious Diseases, Department of Internal Medicine, Emory University School of Medicine, Atlanta, Georgia; and Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Georgia Grady Health System, Atlanta.

**Preeti N. Malani, MD, MSJ** Division of Infectious Diseases, Department of Internal Medicine, University of Michigan, Ann Arbor; and Associate Editor, *JAMA*. [JAMA Patient Page](#)

### Corresponding

**Author:** Carlos del Rio, MD, Emory University School of Medicine, 49 Jesse Hill Jr Dr, FOB Room 201, Atlanta, GA 30303 ([cdelrio@emory.edu](mailto:cdelrio@emory.edu)).

**Now in the third year** of the coronavirus pandemic, well after the Omicron variant surge, both in the US and globally the number of daily cases had been declining to their lowest levels in more than 6 months. While it seemed that SARS-CoV-2 was moving toward endemicity, US infections are again rising in May 2022, and the reported number of cases is likely a gross underestimate of actual infections because many infections are unreported with increasing at-home testing. Several factors help explain the current trends: The emergence of the BA.2 subvariant of

Omicron and the more recently identified subvariant called BA.2.12.1, the limited durability of protection from infection both from vaccination and prior infection, and lifting of mandates (such as mask use) and other restrictions across the country.

### Subvariants of Omicron

After the Omicron variant (BA.1) was first identified in South Africa in November 2021, it quickly spread worldwide, outcompeting other variants and rapidly becoming the pre-dominant variant in many countries including the US.<sup>1</sup> Since then, several lineages and sublineages have emerged. The most common now are BA.1, BA.1.1, BA.2, and BA.2.12.1.

The BA.2 variant effective reproduction number is 1.4-fold higher than that of BA.1.<sup>2</sup> Its viral fitness is due to 53 mutations, 29 of which are in the spike protein. The clinical manifestations of BA.2 infection are similar to those of BA.1, with mild upper respiratory symptoms (such as sore throat and pharyngitis) being common. In addition, many patients report gastrointestinal symptoms (eg, diarrhea, nausea, and vomiting) along with nonspecific symptoms (eg, muscle aches, headache, nasal congestion, and fatigue).

First identified in New York, BA.2.12.1 is now the dominant variant in the US. As of May 25, 2022, about 58% of SARS-CoV-2 isolates sequenced were BA.2.12.1. This subvariant has additional spike mutations S704L and L452Q on top of BA.2 background. The L452Q mutation had been previously observed in the Delta and Lambda variants and allows the virus to attach more tightly to the angiotensin-converting enzyme 2 receptor and thus become more transmissible.<sup>3</sup> Prior infection with BA.1 appears to offer minimal cross-immunity to BA.2.12.1, thus people who had Omicron infection with BA.1 can also get infected with BA.2.12.1.<sup>4</sup>

Two other variants, BA.4 and BA.5, have recently emerged in South Africa and in Europe and have been identified as Variants of Concern Lineages Under Monitoring by

the World Health Organization. Like other Omicron subvariants, BA.4 and BA.5 appear to be significantly more transmissible than previously identified (pre-Omicron) variants. While experiments suggest that at least some of the competitive advantage of BA.4 and BA.5 could be due to differences in viral replication, other factors such as immune evasion or intrinsic transmissibility could also contribute to the observed differences in epidemic growth rates. As with BA.2.12.1, individuals previously infected with an earlier Omicron variant (BA.1) do not appear to be well protected against infection from BA.4 and BA.5. Fortunately, BA.4 and BA.5 do not seem to cause more severe disease than previous variants. In the US, BA.2 and BA.2.12.1 remain the dominant variants as of mid May 2022, but it is likely that more cases will result from BA.4 and BA.5 infection as the summer approaches and as more cases are imported from South Africa and Europe. In summary, since the emergence of Omicron, SARS-CoV-2 is rapidly becoming much more efficient in transmission and more likely to evade immunity.

### Controlling Spread of COVID-19

As the initial Omicron wave was receding in the US, the Centers for Disease Control and Prevention (CDC) launched on March 3, 2022, the COVID-19 Community Level indicator, a new tool to help communities decide what prevention steps to take. Levels can be low, medium, or high as determined by hospital bed utilization, hospital admissions, and total number of COVID-19 cases in a community. When the tool was first launched, more than 90% of the US population lived in locations with a low or medium COVID-19 Community Level where indoor masking was not considered essential. As of May 25, 2022, when daily reported COVID-19 cases are increasing and approaching 100,000, about 9.2% of counties are in the high Community Level.<sup>5</sup> This contrasts with the CDC Community Transmission indicator in which more than two-thirds of counties in the US are in the high transmission level. The existence of 2 different CDC metrics

that provide almost opposing information and longer require masking, people who wish 5 masks available in the US were found to con- sider masks are left wondering what to do and if to be standard or counterfeit fusion and makes recommendations on the use of 1-way masking (wearing a mask while the during evaluations conducted in 2020. KN95 se of nonpharmacologic interventions, surrounding individuals are not masked) pro- s do not meet National Institute for such as masking, more difficult. In addition, focusing on the COVID-19 Community Levels metric contributes to the myth that the pandemic is over, that 1-way masking is still effective but Occupational Safety and Health (NIOSH) standards. NIOSH-approved respirators that for maximum protection the person should have an approval label on or within the something everyone wants to believe is true but certainly is not. wear a high-quality and well-fitting mask such as for the most part, have headbands instead of ear loops. Given the fact that many places are not a KN95.<sup>6</sup> However, an estimated 60% of KN95

[jama.com](https://jama.com)

JAMA Published online May 27, 2022 E1 © 2022 American Medical Association. All rights reserved.

Downloaded From: <https://jamanetwork.com/> by RUBEN PIACENTINI on 06/10/2022

Opinion [Viewpoint](#)

### COVID-19 Vaccine Boosters

Mutations present in Omicron and its subvariants have been associated not only with increased transmissibility, but also with immune evasion to both vaccination and immunity after infection. Primary immunization with 2 doses of mRNA vaccines provides limited protection against symptomatic disease caused by Omicron that increases substantially after a booster.<sup>7</sup> mRNA vaccines offer a similar degree of protection against BA.1 and BA.2 although protection against infection and symptomatic disease wanes within months of a third dose.<sup>8</sup> In contrast, vaccine-associated protection against severe illness, including hospitalization and death, remains durable.

Given the decline in protection against symptomatic infection, a fourth dose of mRNA vaccine in high-risk individuals (older adults and those who are highly immunocompromised) has been proposed as a preventive strategy. Data from Israel suggest that a fourth dose is associated with lower risk of symptomatic infection, hospitalization, severe disease, and death among older adults, but has only marginal benefit in most individuals for whom protection against severe disease remains after a primary series and third dose.<sup>9</sup> Currently, a fourth vaccine dose is recommended by the CDC only for individuals older than 50 years who received their booster at least 4 months ago and those 12 years and older who are moderately or severely immunocompromised and who have gotten their booster at least 4 months prior. As is the case with other vaccines, the goal of vaccination against COVID-19 is to protect against serious illness, not prevent all infections. However, it is increasingly clear that with Omicron and its many subvariants, protection from severe disease requires 3 doses of an mRNA vaccine, something the CDC has termed “up-to-date.” However, less than 50% of those older than 12 years and only 69% of those 65 years and older have received a third vaccine dose.

Emerging data suggest that prior COVID-19 infection provided robust protection against the Alpha, Beta, and Delta variants but limited protection against Omicron. However, the combination of a previous SARS-CoV-2 infection and vaccination (“hybrid immunity”), seems to confer the greatest protection against symptomatic infection although clinically, repeat infections are being seen frequently.

### Therapeutics

With the increased availability of effective antiviral agents and monoclonal antibodies for the treatment of COVID-19, the Biden administration launched a new initiative called Test to Treat in March 2022.<sup>10</sup> Yet,

the program has been underutilized because of logistical barriers, such as the need to initiate treatment within 5 days of symptom onset, and practical concerns, such as serious drug-drug interactions, need for dosing according to kidney function, and potential teratogenicity. Making COVID-19 testing readily available along with rapid linkage to treatment resources will require substantial efforts and a sustainable infrastructure. Planning must ensure that these resources are easily available to those individuals who are not connected to the health care system.

In addition, as nirmatrelvir/ritonavir is increasingly being prescribed, reports of relapse, both with recurrence of symptoms and a positive antigen test result, shortly after ending a 5-day course of treatment are being described. As of now, it is unclear why or how often this is occurring, but these observations warrant monitoring at a population level. In addition, it is unclear whether a longer course of treatment (eg, 10 days rather than 5) or an additional course of treatment is indicated after relapse. Of note, the clinical trials that have evaluated both nirmatrelvir/ritonavir and molnupiravir were conducted among vaccinated individuals at high risk for severe illness. In contrast, most people who are recurrently being prescribed oral antivirals have been vaccinated. Thus, there are currently no data on the efficacy or response to treatment among vaccinated individuals; however, until the data become available, it is reasonable to prescribe these medications to people who are infected and at high risk of severe COVID-19.

### Conclusions

While many questions remain about the future of the pandemic, it is clear that SARS-CoV-2 will not be fully eradicated. This means continuing to adapt to life with COVID-19 and recognizing that during the next phase of the pandemic, there will be times when community transmission will be low and precautions can be “dialed down” and times when increased transmission will require mitigation efforts to be “dialed up.”

If COVID-19 moves toward endemicity, then it should not disrupt everyday life. However, with ongoing transmission and with an estimated 10% to 30%

of individuals experiencing long COVID symptoms risk of long COVID and thus continuing to focus after infection, this issue will require careful attention to further on improving vaccination rates must remain define the syndrome and possible intervention (such as the cornerstone of COVID 19 prevention and mitigation not the RECOVER cohort study at the National Institutes of Health). Data suggest that vaccination can decrease the

#### ARTICLE INFORMATION

**Published Online:** May 27, 2022.  
doi:10.1001/jama.2022.9655

**Correction:** This article was corrected online June 1, 2022, to fix incorrect information on KN95 masks in the eighth paragraph.

**Conflict of Interest Disclosures:** Dr del Rio reported serving as co-principal investigator in the NIH/NIAID Emory Vaccine and Treatment Evaluation Unit. No other disclosures were reported.

#### REFERENCES

1. Del Rio C, Omer SB, Malani PN. Winter of Omicron. *JAMA*. 2022;327(4):319-320.
2. Yamasoba D, Kimura I, Nasser H, et al.

Virological characteristics of SARS-CoV-2 BA.2 variant. *bioRxiv*.

Preprint posted February 15, 2022.  
doi:10.1101/2022.02.14.480335

3. Tchesnokova V, Kulasekara H, Larson L, et al. Acquisition of the L452R mutation in the ACE2-binding interface of spike protein triggers recent massive expansion of SARS-CoV-2 variants. *J Clin Microbiol*. 2021;59(11):e0092121.

4. Cao Y, Yisimayi A, Jian F, et al. BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron infection. *bioRxiv*. Preprint posted May 2, 2022. doi:10.1101/2022.04.30.489997

5. Centers for Disease Control and Prevention. COVID-19 Community Levels. Accessed May 23, 2022.  
<https://www.cdc.gov/coronavirus/2019-ncov/science/community-levels.html>

6. Andrejko KL, Pry JM, Myers JF, et al; California COVID-19 Case-Control Study Team. Effectiveness of face mask or

respirator use in indoor public settings for prevention of SARS-CoV-2 infection—California, February–December 2021. *MMWR Morb Mortal Wkly Rep*. 2022;71(6):212-216. 7. Accorsi EK, Britton A, Fleming-Dutra KE, et al. Association between 3 doses of mRNA COVID-19 vaccine and symptomatic infection caused by the SARS-CoV-2 Omicron and Delta variants. *JAMA*. 2022;327(7):639-651.

8. Lassaunière R, Polacek C, Frische A, et al. Neutralizing antibodies against the SARS-CoV-2 Omicron variant (BA.1) 1 to 18 weeks after the second and third doses of the BNT162b2 mRNA vaccine. *JAMA Netw Open*. 2022;5(5):e2212073. 9. Bar-On YM, Goldberg Y, Mandel M, et al. Protection by a fourth dose of BNT162b2 against Omicron in Israel. *N Engl J Med*. 2022;386(18):1712-1720.

10. Gandhi RT, Malani PN, Del Rio C. COVID-19 therapeutics for nonhospitalized patients. *JAMA*. 2022;327(7):617-618.